

Self-Stark Limit on Brightness Temperature in Cosmic Masers

V. I. SLYSH

Arecibo Observatory, HC3 Box 53995, Arecibo PR 00612

Astro Space Center, Profsoyuznaya st. 84/32, 117997 Moscow, Russia

Abstract. The radiation broadening of narrow maser lines by its own emission which can also be described as resonance Self-Stark effect puts an upper limit on the brightness temperature of cosmic masers. It is shown that available measurements of the brightness temperature of OH and water masers are consistent with the observed narrow line widths only if the maser emission is highly directive.

1. Introduction

Methanol, OH and water masers are generally seen toward star-forming regions and exhibit narrow, strong emission line features coming from bright small spots. The brightness temperature T_b of the spots is very high, from 10^9 K for methanol masers, to 6×10^{12} K for OH masers, and 10^{15} K for water masers. In many cases the above values are only lower limits since maser spots are barely resolved with VLBI. Variability time scale often indicates, in an indirect way, a smaller linear size and still higher brightness temperature. There must be physical limits on the brightness temperature as there are for thermal and synchrotron emission in continuum sources. The brightness temperature of the thermal free-free emission from ionized gas depends on the kinetic temperature T_e of electrons and optical depth. It is the highest for optically thick sources and is equal to the kinetic temperature. In H II regions the kinetic temperature of electrons is limited to $T_e = 10,000$ K by the energy losses from electrons through the line emission of heavy elements. Similarly, in synchrotron sources the brightness temperature is at maximum in optically thick sources, and is equal to the kinetic temperature of relativistic electrons. The kinetic energy of relativistic electrons is limited to 10^{11} – 10^{12} K by the energy loss through inverse Compton emission on synchrotron photons (Kellermann & Pauliny-Toth 1969). The same arguments can be applied to molecular line emission: in the optically thick sources the brightness temperature in the line is limited by the gas kinetic temperature. The molecular maser sources make an exception because the absorption coefficient and optical depth are both negative due to the population inversion: the brightness temperature of the maser line emission may increase indefinitely with the optical depth. However other mechanisms can limit the brightness temperature of the maser lines, such as the radiation line broadening, or the resonance Stark-effect caused by the maser radiation field (we call it Self-Stark limit by analogy with Self-Compton limit).

2. Radiation, or resonance Stark broadening of maser lines

The natural band width of a molecular emission line in the absence of external field is $\Delta\nu = A/2\pi$, where $A = 64\pi^4\nu_0^3\mu^2/3hc^3$ is spontaneous emission rate at the frequency of the line ν_0 , μ is the dipole-moment matrix element for the transition, and h is Planck constant. This is a very small band width compared to the observed line widths. For OH line at 1665 MHz, $A = 7.1 \times 10^{-11} \text{ s}^{-1}$, and for water line at 22.235 GHz, $A = 2 \times 10^{-9} \text{ s}^{-1}$, and the natural band widths are 10^{-11} Hz and $3 \times 10^{-10} \text{ Hz}$, respectively. In strong masers like W3(OH) or W49 every OH or H_2O molecule is exposed to the intense radiation field, produced by all other molecules in the maser emission region. That causes frequent up and down transitions between the energy levels involved, due to induced emission and absorption. The time which a molecule spends on the energy level is greatly reduced, by a factor $2kT_b/h\nu_0$. The linewidth is increased by the same factor which can be very large when the brightness temperature is high. For OH maser with brightness temperature $6 \times 10^{12} \text{ K}$ the linewidth is $1,700 \text{ Hz}$ (0.3 km s^{-1}), and for water maser with the brightness temperature 10^{15} K it is 0.6 MHz (8 km s^{-1}). This linewidth is comparable to or even greater than observed line widths in masers. Einstein induced emission and absorption coefficients can be used only in the weak field case. For the strong field which is the case in water masers a correct description of the interaction between a molecule and the narrow band radiation field is the resonance Stark effect approach. In the weak field the resonance Stark effect is quadratic with respect to the electric field strength E , and causes the line broadening. In the strong field the resonance Stark effect becomes linear, and causes line splitting into a main line at the transition frequency ν_0 and two satellite lines at frequencies $\nu_0 \pm \mu E/h$ for non-degenerate energy levels such as are levels of the OH line transition at 1665 MHz. The energy levels of the H_2O transition at 22.235 GHz are degenerate, and are split in several satellite components (Slysh 1973). AC electric field strength E is related to the brightness temperature T_b of the maser radiation of band width $\delta\nu$ as $E = 8\pi(kT_b)^{0.5}\nu_0(\delta\nu)^{0.5}c^{-1.5}$. This relation holds for the isotropic distribution of the radiation intensity. If the radiation is concentrated in the solid angle Ω , the electric field strength must be multiplied by $(\Omega/4\pi)^{0.5}$, with the corresponding reduction of the line splitting. For the line broadening the reduction factor is $\Omega/4\pi$.

3. Brightness temperature of H_2O and OH masers

The strongest spectral feature in H_2O maser W3OH is completely unresolved on all VLBA baselines (Fig. 1), and its brightness temperature exceeds $4 \times 10^{15} \text{ K}$. Another strong H_2O maser, an outburst in Orion-KL, was measured with the space radio telescope *HALCA* to have brightness temperature $T_b = 10^{16} \text{ K}$ (Kobayashi et al 2000). OH maser OH 34.26+0.15 was unresolved on baselines from ground to space radio telescope *HALCA*, and the brightness temperature exceeds $4 \times 10^{12} \text{ K}$ (Fig. 2).




Figure 1. Correlated flux vs baseline for the strongest H_2O maser spectral feature in W3OH obtained from VLBA archive data

4. Directivity of the maser emission

No line splitting due to the resonance Stark effect was observed in H_2O or OH masers. Also, no significant radiation line broadening is seen in the line profiles. This means that the electric field strength in the masers is weak enough in spite of the very high brightness temperature, because of the small solid angle in which emission is directed. Assuming that the radiation broadening contributes to the total linewidth less than $1/5$ of the linewidth, one can put an upper limit on the solid angle. One has for OH maser OH 34.26+0.15 an upper limit for the emission solid angle $\Omega/4\pi=0.02$. For H_2O masers in W3OH one has $\Omega/4\pi=0.005$, and for the outburst H_2O maser in Orion-KL $\Omega/4\pi=0.002$ as the upper limits.

5. Conclusion

The brightness temperature of cosmic masers is limited by the radiation broadening in the intense radiation field at the frequency of the maser emission. Up




Figure 2. A map of the most compact OH-maser emission spectral feature in OH 34.26+0.15. Participating telescopes were phased VLA, 70 m Tidbinbilla and 8 m space telescope *HALCA*. Synthesized beam was 2.4×1.1 mas

to now no radiation broadening has been observed in OH and H₂O masers, which can be due to the very small solid angle in which the emission is directed. The theory of the resonance radiation broadening allows us to put stringent upper limits on the emission solid angle. This may be important for the further development of cosmic maser models.

References

- Kellermann, K. I. & Pauliny-Toth, I. I. K. 1969, ApJ, 155, L71
 Kobayashi, H., Shimoikura, T., Omodaka, T. & Diamond, P.J. 2000 in Astrophysical Phenomena Revealed by Space VLBI, ed. H. Hirabayashi, P. G. Edwards & D. W. Murphy, 109
 Slysh, V. I. 1973, Astrophysical Letters, 14, 213